

Gravitino condensation in fivebrane backgrounds

Noriaki Kitazawa *

Department of Physics, Tokyo Metropolitan University, Hachioji, Tokyo 192-0397, Japan
(February 1, 2008)

Abstract

We calculate the tension of the D3-brane in the fivebrane background which is described by the exactly solvable $SU(2)_k \times U(1)$ world-sheet conformal field theory with large Kač-Moody level k . The D3-brane tension is extracted from the amplitude of one closed string exchange between two parallel D3-branes, and the amplitude is calculated by utilizing the open-closed string duality. The tension of the D3-brane in the background does not coincide with the one in the flat space-time even in the flat space-time limit: $k \rightarrow \infty$. The finite curvature effect should vanish in the flat space-time limit and only the topological effect can remain. Therefore, the deviation suggests the condensation of gravitino and/or dilatino which has been expected in the fivebrane background as a gravitational instanton.

Typeset using REVTeX

*e-mail: kitazawa@phys.metro-u.ac.jp

I. INTRODUCTION

It has been pointed out that the local supersymmetry can be spontaneously broken by the gravitino condensation due to the topological effect of the quantum gravity [1,2]. An explicit calculation in the framework of the Euclidean quantum gravity theory is given in Ref. [2], and the finite value of the gravitino pair condensate $\langle \psi_{\mu\nu} \psi_{\mu\nu} \rangle$ is obtained, where $\psi_{\mu\nu} = D_\mu \psi_\nu - D_\nu \psi_\mu$ is the field strength of the gravitino field ψ_μ and D_μ is the covariant derivative. This mechanism is very attractive, since it is less model-dependent.

It is claimed that this mechanism might work also in the low energy effective theory of the superstring theory. The existence of the appropriate zero modes for the gravitino pair condensation in the fivebrane background is explicitly shown in Ref. [3] in the framework of the Euclidean quantum gravity theory. Since the fivebrane background is exactly described by using the world-sheet conformal field theory [4–6] as a solution of the string theory, it is interesting to confirm this suggestion in the string theory from the first principle.

A D-brane act as a source for closed strings. The D-brane tension is nothing but the coupling strength between a D-brane and a single lowest-lying closed string state. In the low-energy limit (considering only the lowest-lying (massless) states of the string excitation) a single Dp-brane in the type II string theory is described by the effective action

$$S_p = -T_p \int d^{p+1} \xi \left\{ e^{-\Phi} [-\det (G_{ab} + B_{ab} + 2\pi\alpha' F_{ab})]^{1/2} + \mathcal{O}(\alpha') \right\}, \quad (1)$$

where T_p is the tension of the Dp-brane, ξ^a is the Dp-brane coordinate ($a = 0, 1, \dots, p$), Φ is the dilaton field, G_{ab} is the pull back of the string metric $G_{\mu\nu}$ ($\mu, \nu = 0, 1, \dots, 9$), B_{ab} is the pull back of the B-field $B_{\mu\nu}$, and F_{ab} is the gauge field from the open string in the Dp-brane. (Here, we neglect the contribution from R-R fields.) After rescaling the string metric to the Einstein metric $g_{\mu\nu} = e^{-\phi/2} G_{\mu\nu}$ with $\Phi = \langle \Phi \rangle + \phi$, the couplings of a single dilaton or graviton to a Dp-brane can be obtained. The coupling is proportional to the effective Dp-brane tension $\tau_p = T_p e^{-\langle \Phi \rangle}$.

In case of the curved space-time background the higher order terms in Eq.(1) contribute to the effective tension. The higher order R^2 terms, for example, are obtained in Ref. [8] in case of D3-brane:

$$S_3^{R^2} = -T_3 \frac{(2\pi\alpha')^2}{96} \int d^4 \xi e^{-\Phi} \text{tr} (R \wedge *R) + \dots, \quad (2)$$

where $R^c_d = \frac{1}{2} R^c_{dab} d\xi^a \wedge d\xi^b$ and $*R^c_d = \frac{1}{4} \sqrt{-G} \epsilon_{ab}{}^{a'b'} R^c_{da'b'} d\xi^a \wedge d\xi^b$. The non-trivial background gives rise to a vacuum expectation value of the curvature tensor, and the effective D-brane tension is modified. The supersymmetric counter part of the effective action of Eq.(2) can be obtained by using the superfield method described in Ref. [9–12]. We find

$$S_3^{\psi^2} \propto T_3 \frac{(2\pi\alpha')^2}{96} \int d^4 \xi \sqrt{-G} e^{-\Phi} (\partial^c \lambda \partial_c \lambda) (\psi^{ab} \psi_{ab}) + \dots, \quad (3)$$

where ψ is a linear combination of two gravitinos corresponding to $N = 2$ supersymmetry of the type II theory, and λ , which has mass dimension $-1/2$, is a Goldstone fermion associated with the half supersymmetry breaking by the D3-brane. The vacuum expectation value of

composite operators, like $\langle(\partial^c\lambda\partial_c\lambda)(\psi^{ab}\psi_{ab})\rangle$ in this example, also modify the effective D-brane tension. Since the low-energy amplitude of one closed string exchange between two parallel D-branes (Fig.1) is determined by the effective D-brane tension, the non-trivial background effect may modify the low-energy amplitude.

In this paper we calculate the amplitude of Fig.1 in the fivebrane background, and compare it with the same amplitude in the flat space-time. Since the superstring theory in the fivebrane background, $M^6 \otimes W_k^{(4)}$, where M^6 is the six-dimensional flat Minkowski space-time and $W_k^{(4)} \equiv \text{SU}(2)_k \times \text{U}(1)$ is the four-dimensional non-trivial space, can be exactly constructed, we can calculate the amplitude exactly. In the flat space-time limit, $k \rightarrow \infty$, the finite curvature effect disappears in the amplitude, and only the topological effect should remain. (Notice also that the space-time background interpretation of this string solution exists only for large Kač-Moody level k .) We will see in the low-energy limit that the amplitude in the flat space-time limit does not coincide with the one in the flat space-time. Namely, the effective D-brane tension in the flat space-time limit does not coincide with the one in the flat space-time. This suggests the gravitino and/or dilatino condensation by the topological effect in the string theory.

In the next section we review the string solution in the fivebrane background. In section III the amplitude of Fig.1 is calculated both in the flat space-time and in the fivebrane background. The flat space-time limit of the amplitude is discussed. It will be found that the effective D-brane tension in the flat limit does not coincide with the one in the flat space-time. In the last section we summarize the result.

II. STRING THEORY ON FIVEBRANE BACKGROUNDS

In this section we briefly review the exact string solution in the fivebrane background (see Refs. [4,5] for details). We concentrate on the construction of the partition functions of closed and open strings in the fivebrane background $M^6 \otimes W_k^{(4)}$.

In case of the flat space-time the world-sheet theory consists of a free bosonic field $X^\mu(z, \bar{z})$ and two free fermionic fields $\psi^\mu(z)$ and $\tilde{\psi}^\mu(\bar{z})$. The system has $(N, \tilde{N}) = (1, 1)$ superconformal symmetry with the central charge $c_m = \tilde{c}_m = 10 + 10/2 = 15$ which is cancelled by the ghost contribution $c_g = \tilde{c}_g = -26 + 11 = -15$. The non-trivial background $M^6 \otimes W_k^{(4)}$ can be described by replacing unconstrained fields X^μ with $\mu = 6, 7, 8, 9$ to the fields constrained on the group manifold $W_k^{(4)} = \text{SU}(2)_k \times \text{U}(1)$. Namely, a part of the world-sheet theory corresponding to the space-time dimensions of $\mu = 6, 7, 8, 9$ is replaced by the $\text{SU}(2)_k \times \text{U}(1)$ WZW model with $(N, \tilde{N}) = (4, 4)$ superconformal symmetry which has the same central charge of the replaced part $c_{WZW} = 6$.

The holomorphic part of the $\text{SU}(2)_k \times \text{U}(1)$ WZW model is described by three $\text{SU}(2)_k$ bosonic currents $J_i(z)$ ($i = 1, 2, 3$), one free bosonic current $J_4(z) = \partial X^{\mu=6}$ and four free fermionic fields Ψ_a ($a = 1, 2, 3, 4$). These currents and fields satisfy the following the operator product expansion.

$$J_i(z)J_j(z') \sim -\frac{k}{2} \frac{\delta_{ij}}{(z-z')^2} + \epsilon_{ijl} \frac{J_l}{z-z'}, \quad (4)$$

$$J_4(z)J_4(z') \sim -\frac{1}{(z-z')^2}, \quad (5)$$

$$\Psi_a(z)\Psi_b(z') \sim -\frac{\delta_{ab}}{z-z'}, \quad (6)$$

where k is the Kač-Moody level of $SU(2)_k$. The $N = 4$ superconformal symmetry transformation is generated by the following energy-momentum tensor $T(z)$, supercurrents $G_a(z)$ and $SU(2)_n$ currents $S_i(z)$.

$$T = -\frac{1}{2} \left(\frac{2}{k+2} J_i^2 + J_4^2 - \Psi_a \partial \Psi_a + Q \partial J_4 \right), \quad (7)$$

$$G_i = \sqrt{\frac{2}{k+2}} \left(J_i \Psi_4 - \epsilon_{ijl} J_j \Psi_l + \frac{1}{2} \epsilon_{ijl} \Psi_4 \Psi_j \Psi_l \right) - J_4 \Psi_i - Q \partial \Psi_i, \quad (8)$$

$$G_4 = \sqrt{\frac{2}{k+2}} \left(J_i \Psi_i + \frac{1}{3!} \epsilon_{ijl} \Psi_i \Psi_j \Psi_l \right) + J_4 \Psi_4 + Q \partial \Psi_4, \quad (9)$$

$$S_i = \frac{1}{2} \left(\Psi_4 \Psi_i + \frac{1}{2} \epsilon_{ijl} \Psi_j \Psi_l \right). \quad (10)$$

The background charge Q determines the gradient of the linear-dilaton background $\Phi = QX^{\mu=6}$. The world-sheet field $X^{\mu=6}(z, \bar{z})$ is called Feigin-Fuchs field. The value $Q = \sqrt{2/(k+2)}$ is required for the correct central charge of $c_{WZW} = 6$, and the Kač-Moody level n of $SU(2)_n$ is fixed to unity. (In the present discussion we set $\alpha' = 2$. For example, $Q = \sqrt{\alpha'/(k+2)} = \sqrt{2/(k+2)}$.) The anti-holomorphic part has exactly the same structure.

A. The closed string partition function

The partition function of the closed string is defined by

$$Z^c(\tau) = \text{Tr} \left[e^{2\pi i \tau_1 P - 2\pi \tau_2 H} \right] = (q\bar{q})^{-d/24} \text{Tr} \left[q^{L_0} \bar{q}^{\tilde{L}_0} \right], \quad (11)$$

where $P = L_0 - \tilde{L}_0$ and $H = L_0 + \tilde{L}_0 - d/24$ are the world-sheet momentum and Hamiltonian operators, $\tau = \tau_1 + i\tau_2$, $q = e^{2\pi i \tau}$, $d = (c + \tilde{c})/2$ with total central charges c and $\tilde{c}$ and L_0 and $\tilde{L}_0$ are Virasoro generators. The torus vacuum amplitude is obtained by integrating this partition function over the moduli parameter τ

$$A_{T^2} = \int_F \frac{d^2 \tau}{4\tau_2} Z^c(\tau), \quad (12)$$

where F denotes the fundamental region of the moduli space. The partition function can be decomposed in two parts $Z^c = Z_B^c \times Z_F^c$, where Z_B^c is the bosonic (X^μ and bc -ghosts) contribution and Z_F^c is the fermionic (ψ^μ , $\bar{\psi}^\mu$ and $\beta\gamma$ -ghosts) contribution.

The construction of the partition function in fivebrane backgrounds can be understood by starting with the partition function in the flat space-time. In case of $D = 10$ dimensional flat space-time the bosonic contribution Z_B^c is obtained as

$$Z_B^c(\tau) = \frac{iV_D}{4\pi^2\alpha'\tau_2} (Z_X^c(\tau))^{D-2}, \quad (13)$$

where

$$Z_X^c(\tau) = (4\pi^2\alpha'\tau_2)^{-1/2} |\eta(\tau)|^{-2} \quad (14)$$

is the contribution of one space-time dimension. The factor i in Eq.(13) originates from the negative sign of the time component of the flat space-time Minkowski metric. The fermionic contribution is obtained as

$$Z_F^c(\tau) = Z_\psi(\tau) (Z_\psi(\tau))^*, \quad (15)$$

where

$$Z_\psi(\tau) = \frac{1}{2} \sum_{\alpha,\beta} (-1)^{\alpha+\beta-\alpha\beta} \left(\frac{\theta_{\beta}^{(\alpha)}(\tau)}{\eta(\tau)} \right)^{(D-2)/2} \quad (16)$$

for the type IIB theory. The two sectors, Neveu-Schwarz and Ramond sectors, are included by the summation over $\alpha = 0, 1$. The GSO projection is realized by the summation over $\beta = 0, 1$.

We modify the above result to the case of the fivebrane background $M^6 \otimes W_k^{(4)}$. The bosonic contribution of the M^6 part can be obtained by setting $D = 6$ in Eq.(13). The contribution of the Feigin-Fuchs field (U(1) part of $W_k^{(4)}$) is given by Eq.(14) times V_1 as the contribution of a single space-like coordinate, if the state of momentum k is defined as

$$|k\rangle =: e^{i(k+i\frac{Q}{\alpha'})X^{\mu=6}(0,0)} : |0\rangle \quad (17)$$

corresponding to the modification of L_0 and $\tilde{L}_0$. By the effect of the Feigin-Fuchs field, the whole mass spectrum is uniformly lifted up by $\Delta m^2 = (Q/\alpha')^2$, and there is no massless state.

The bosonic contribution of the $SU(2)_k$ part should be described by the Virasoro character

$$\chi_k^n(\tau) = q^{s_{n,k}} \text{ch}_{n,k}(\tau), \quad (18)$$

where $n = 0, 1, \dots$ denotes the representation of $SU(2)$, $s_{n,k} = \frac{((n+1)/2)^2}{k+2} - \frac{3}{24}$ and $\text{ch}_{n,k}(\tau)$ is the $SU(2)_k$ character which corresponds to $\text{Tr}(q^{L_0})$ (see Refs. [13,14] for precise definitions). The actual form of the Virasoro character is

$$\chi_k^n(\tau) = \frac{1}{(\eta(\tau))^3} \cdot 2(k+2) \sum_{m=-\infty}^{\infty} \left(m + \frac{n+1}{2(k+2)} \right) e^{2\pi i(k+2)\tau \left(m + \frac{n+1}{2(k+2)} \right)^2}. \quad (19)$$

The Virasoro character with $n \leq k$ is a unitary representation under the modular transformation:

$$\chi_k^n(-1/\tau) = \sum_{n'=0}^k S_{nn'} \chi_k^{n'}(\tau), \quad S_{nn'} = \left(\frac{2}{k+2} \right)^{1/2} \sin \left(\pi \frac{(n+1)(n'+1)}{k+2} \right), \quad (20)$$

$$\chi_k^n(\tau+1) = \sum_{n'=0}^k T_{nn'} \chi_k^{n'}(\tau), \quad T_{nn'} = \delta_{nn'} e^{i\frac{\pi}{2} \left(\frac{(n+1)^2}{k+2} - \frac{1}{2} \right)}. \quad (21)$$

Therefore, a combination

$$\sum_{n=0}^k \chi_k^n(\tau) (\chi_k^n(\tau))^* \quad (22)$$

is a modular invariant, but this is not appropriate for the $SU(2)_k$ contribution to the partition function, since the whole partition function (or the vacuum amplitude) should be modular invariant.

The modular invariance and the extraction of the physical state by the GSO projection lead the following partition function for even k .

$$\begin{aligned} Z^c(\tau) &= \frac{iV_6}{4\pi^2\alpha'\tau_2} (Z_X^c(\tau))^4 \cdot V_1 Z_X^c(\tau) \\ &\times \sum_{\alpha,\beta} \sum_{\bar{\alpha},\bar{\beta}} \sum_{\gamma,\delta} \frac{1}{2} (-1)^{\alpha+\beta-\alpha\beta} \left(\frac{\theta\left(\frac{\alpha}{\beta}\right)(\tau)}{\eta(\tau)} \right)^2 \left(\frac{\theta\left(\frac{\alpha+\gamma}{\beta+\delta}\right)(\tau)}{\eta(\tau)} \right)^2 \\ &\times \frac{1}{2} (-1)^{\bar{\alpha}+\bar{\beta}-\bar{\alpha}\bar{\beta}} \left(\left(\frac{\theta\left(\frac{\bar{\alpha}}{\bar{\beta}}\right)(\tau)}{\eta(\tau)} \right)^2 \left(\frac{\theta\left(\frac{\bar{\alpha}+\gamma}{\bar{\beta}+\delta}\right)(\tau)}{\eta(\tau)} \right)^2 \right)^* \\ &\times \frac{1}{2} (-1)^{\delta(\alpha+\bar{\alpha}+\gamma k/2)} Z_k \left[\begin{smallmatrix} \gamma \\ \delta \end{smallmatrix} \right](\tau), \end{aligned} \quad (23)$$

where

$$Z_k \left[\begin{smallmatrix} \gamma \\ \delta \end{smallmatrix} \right](\tau) = \sum_{n=0}^k e^{i\pi\delta n} \chi_k^n(\tau) (\chi_k^{n+\gamma(k-2n)}(\tau))^* \quad (24)$$

which covariantly transforms under the modular transformation as

$$Z_k \left[\begin{smallmatrix} \gamma \\ \delta \end{smallmatrix} \right](-1/\tau) = e^{i\pi\gamma\delta k} Z_k \left[\begin{smallmatrix} \delta \\ \gamma \end{smallmatrix} \right](\tau), \quad (25)$$

$$Z_k \left[\begin{smallmatrix} \gamma \\ \delta \end{smallmatrix} \right](\tau+1) = e^{-i\pi\gamma^2 k/2} Z_k \left[\begin{smallmatrix} \gamma \\ \delta + \gamma \end{smallmatrix} \right](\tau). \quad (26)$$

In Eq.(23) the summation over $\delta = 0, 1$ realizes the additional GSO projection which reduces the space-time supersymmetry by a factor of 2. This corresponds to the fact that the fivebrane configuration is a BPS saturated state which preserves only half of the supersymmetry of the theory. The fermionic contribution is the same in case of the flat space-time, except for the modification of the contribution from the fermions in $W_k^{(4)}$ by the additional GSO projection. The summation over $\gamma = 0, 1$ is required by the modular invariance, and a twisted sector ($\gamma = 1$) should exist.

We discuss the flat space-time limit $k \rightarrow \infty$ of the partition function of Eq.(23). Consider first the $k \rightarrow \infty$ limit of the Virasoro character of Eq.(19). Since $\tau_2 \geq 1$ in the fundamental region of the moduli space, there is an exponential suppression factor in the summation over m . Considering $(n+1)/2(k+2) < 1/2$, a contribution with $m=0$ dominates in $k \rightarrow \infty$ limit. Therefore,

$$\lim_{k \rightarrow \infty} \chi_k^n(\tau) = \frac{1}{(\eta(\tau))^3} \cdot (n+1) e^{2\pi i(k+2)\tau \left(\frac{n+1}{2(k+2)}\right)^2}. \quad (27)$$

Then,

$$\lim_{k \rightarrow \infty} Z_k \begin{bmatrix} 0 \\ 0 \end{bmatrix}(\tau) = \frac{1}{|\eta(\tau)|^{-6}} \lim_{k \rightarrow \infty} \sum_{n=0}^k 2(k+2) \left(\frac{n+1}{\sqrt{2(k+2)}} \right)^2 e^{-2\pi\tau_2 \left(\frac{n+1}{\sqrt{2(k+2)}}\right)^2}. \quad (28)$$

We can replace the summation to the integration over the valuable $x = (n+1)/\sqrt{2(k+2)}$:

$$\begin{aligned} \lim_{k \rightarrow \infty} Z_k \begin{bmatrix} 0 \\ 0 \end{bmatrix}(\tau) &= \frac{1}{|\eta(\tau)|^{-6}} \lim_{k \rightarrow \infty} 2(k+2) \int_0^{\frac{k}{\sqrt{2(k+2)}}} \frac{dx}{1/\sqrt{2(k+2)}} x^2 e^{-2\pi\tau_2 x^2} \\ &= \frac{1}{|\eta(\tau)|^{-6}} \lim_{k \rightarrow \infty} (2(k+2))^{3/2} \int_0^\infty dx x^2 e^{-2\pi\tau_2 x^2} \\ &= (Z_X(\tau))^3 \lim_{k \rightarrow \infty} 2\pi^2 \left(\frac{2}{Q} \right)^3, \end{aligned} \quad (29)$$

where we use $\alpha' = 2$ in the closed string Virasoro character due to the relation between mode operators and the space-time momentum operator: $\alpha_0^\mu = \tilde{\alpha}_0^\mu = \sqrt{\alpha'/2} p^\mu$. The factor $2\pi^2(2/Q)^3$ is the volume of S^3 of the radius $2/Q$, which is the volume of the $SU(2)_k$ manifold. Since $Q \rightarrow 0$ in $k \rightarrow \infty$ limit, the factor becomes V_3 in the flat space-time limit, and Eq.(29) becomes just the contribution of three space-like coordinates in the flat space-time. We can show in the same way that $Z_k \begin{bmatrix} \gamma \\ \delta \end{bmatrix}(\tau)$ with other values of γ and δ vanishes in $k \rightarrow \infty$ limit. Therefore, we see that the partition function of Eq.(23) reduces to the one in the flat space-time with an additional $1/2$ factor in the flat space-time limit $k \rightarrow \infty$. This additional factor is the remnant of the additional GSO projection. The supersymmetry which is broken by the fivebrane background does not recover in the simple flat space-time limit. This suggests that the supersymmetry is broken by some topological effects.

B. The open string partition function

The argument which is similar to the closed string is also possible for the open string. The partition function of the (oriented) open string is defined by

$$Z^o(t) = \text{Tr} \left[e^{-2\pi t H} \right] = q^{-c/24} \text{Tr} \left[q^{L_0} \right], \quad (30)$$

where $H = L_0 - c/24$ is the world-sheet Hamiltonian, $q = e^{2\pi i \tau}$ with $\tau = it$ and we use the doubling trick. The cylinder vacuum amplitude is obtained by integrating this partition function over t .

$$A_{C^2} = \int_0^\infty \frac{dt}{2t} Z^o(t). \quad (31)$$

The partition function can be separated in two parts: $Z^o(t) = Z_B^o(t) \times Z_F^o(t)$. In case of the $D = 10$ dimensional flat space-time the bosonic (X^μ and bc -ghosts) contribution is

$$Z_B^o(t) = \frac{iV_D}{8\pi^2\alpha't} (Z_X(t))^{D-2} \quad (32)$$

with

$$Z_X^o(t) = (8\pi^2\alpha't)^{-1/2} (\eta(it))^{-1}. \quad (33)$$

The fermionic (ψ^μ , $\tilde{\psi}^\mu$ and $\beta\gamma$ -ghosts) contribution is

$$Z_F^o(t) = Z_\psi(it), \quad (34)$$

where $Z_\psi(\tau)$ is given in Eq.(16). The Chan-Paton degrees of freedom can be included by multiplying a factor n_{CP}^2 .

We modify the partition function in the flat space-time and obtain the one in the fivebrane background $M^6 \otimes W_k^{(4)}$. The bosonic contribution of the M^6 part is Eq.(32) with $D = 6$. The contribution of the Feigin-Fuchs field is Eq.(33) times V_1 . The bosonic contribution of the $SU(2)_k$ part should be described by a linear combination of Virasoro characters χ_k^n without their complex conjugate [7]. The partition function in the fivebrane background can be obtained as

$$\begin{aligned} Z^o(t) &= \frac{iV_6}{8\pi^2\alpha't} (Z_X^o(t))^4 \cdot V_1 Z_X^o(t) \\ &\times \frac{1}{2} \sum_{\delta} \sum_{n=0}^k (n+1) (-1)^{\delta n} \chi_k^n(it) \\ &\times \frac{1}{2} \sum_{\alpha, \beta} (-1)^{\alpha+\beta-\alpha\beta} \left(\frac{\theta\left(\frac{\alpha}{\beta}\right)(it)}{\eta(it)} \right)^2 (-1)^{\delta\alpha} \left(\frac{\theta\left(\frac{\alpha}{\beta+\delta}\right)(it)}{\eta(it)} \right)^2, \end{aligned} \quad (35)$$

where we consider the additional GSO projection in the same way in the closed string partition function. The fermionic contribution is the same as the one in the flat space-time, except for the modification by the additional GSO projection. The Chan-Paton degrees of freedom can be included by multiplying a factor n_{CP}^2 .

In the flat space-time limit of $k \rightarrow \infty$ the partition function reduces to the one in the flat space-time, except for an additional $1/2$ factor. Consider the limit of the $SU(2)_k$ bosonic contribution

$$Z_k^\delta(t) = \sum_{n=0}^k (n+1) (-1)^{\delta n} \chi_k^n(it) \quad (36)$$

using Eq.(27) in case of $\delta = 0$:

$$\begin{aligned}
\lim_{k \rightarrow \infty} Z_k^0(t) &= \lim_{k \rightarrow \infty} \sum_{n=0}^k (n+1) \chi_k^n(it) \\
&= \lim_{k \rightarrow \infty} \sum_{n=0}^k (n+1) \cdot \frac{1}{(\eta(it))^3} (n+1) e^{-2\pi(k+2)t \left(\frac{n+1}{2(k+2)}\right)^2} \\
&= \frac{1}{(\eta(it))^3} \lim_{k \rightarrow \infty} \sum_{n=0}^k 2(k+2) \left(\frac{n+1}{\sqrt{2(k+2)}} \right)^2 e^{-\pi t \left(\frac{n+1}{\sqrt{2(k+2)}} \right)^2}.
\end{aligned} \tag{37}$$

The summation can be replaced by the integration over the valuable $x = (n+1)/\sqrt{2(k+2)}$.

$$\begin{aligned}
\lim_{k \rightarrow \infty} Z_k^0(t) &= \frac{1}{(\eta(it))^3} \lim_{k \rightarrow \infty} 2(k+2) \int_0^{\frac{k}{\sqrt{2(k+2)}}} \frac{dx}{1/\sqrt{2(k+2)}} x^2 e^{-\pi t x^2} \\
&= \frac{1}{(\eta(it))^3} \lim_{k \rightarrow \infty} (2(k+2))^{3/2} \int_0^\infty dx x^2 e^{-\pi t x^2} \\
&= (Z_X(t))^3 \lim_{k \rightarrow \infty} 2\pi^2 \left(\frac{2}{Q} \right)^3,
\end{aligned} \tag{38}$$

where we use $\alpha' = 1/2$ in the open string Virasoro character due to the relation between a mode operator and the space-time momentum operator: $\alpha_0^\mu = \sqrt{2\alpha'} p^\mu$. Therefore, $Z_k^0(t)$ reduces to just the contribution of three space-like coordinates in the flat space-time. Since we can show in the same way that the flat limit of $Z_k^1(t)$ vanishes, the total partition function reduces to the one in the flat space-time with an $1/2$ factor. This factor is the remnant of the additional GSO projection. The meaning of this factor is the same of the additional factor in the closed string partition function.

III. ONE CLOSED STRING EXCHANGE AMPLITUDE

In this section we calculate the amplitude of Fig.1 both in the flat space-time and in the fivebrane background. The open string cylinder vacuum amplitude, which has been obtained in the previous section, can be utilized for the calculation under the open-closed string duality.

A. The amplitude in the flat space-time

First, we calculate the amplitude of Fig.1 in the flat space-time. This calculation is described in detail in Ref. [15]. Consider two parallel D3-branes separated with distance y and an (oriented) open string whose ends are fixed in different D3-branes. Since the open string has no space-time momentum in the directions of $\mu = 1, 2, 3, 4, 5, 6$ and it is stretched between two D3-branes, in the world-sheet Hamiltonian a part of the momentum term $\alpha' p^\mu p_\mu$ with $\mu = 1, 2, 3, 4, 5, 6$ is replaced by the tension term $y^2/4\pi^2\alpha'$. Considering this replacement in the calculation of the cylinder vacuum amplitude A_{c2} in the previous section gives the amplitude of Fig.1. We should set the Chan-Paton factor $n_{CP}^2 = 2$ to represent the left and right degrees of freedom of the closed string. The result is

$$A^{M^{10}}(y) = \int_0^\infty \frac{dt}{2t} \frac{1}{V_6} \left(8\pi^2 \alpha' t\right)^{6/2} e^{-\frac{y^2}{2\pi\alpha'} t} \cdot 2Z_B^o(t) Z_F^o(t). \quad (39)$$

The world-sheet time coordinate t for the open string should be changed to the one for the closed string, $s = \pi/t$, by using the modular transformation:

$$\eta(it) = t^{-1/2} \eta(i/t), \quad (40)$$

$$\frac{\theta_{(\beta)}^{(\alpha)}(it)}{\eta(it)} = \frac{\theta_{(\alpha)}^{(\beta)}(i/t)}{\eta(i/t)}. \quad (41)$$

Namely,

$$A^{M^{10}}(y) = \frac{iV_4}{2\pi (8\pi^2 \alpha')^5} \int_0^\infty ds \left(8\pi^2 \alpha' \frac{\pi}{s}\right)^3 e^{-\frac{y^2}{2\pi\alpha'} \frac{\pi}{s}} \times (\eta(is/\pi))^{-8} \left\{ Z_\psi^{NS-NS}(is/\pi) + Z_\psi^{R-R}(is/\pi) \right\}, \quad (42)$$

where

$$Z_\psi^{NS-NS}(is/\pi) = \frac{1}{(\eta(is/\pi))^4} \left\{ \left(\theta \begin{pmatrix} 0 \\ 0 \end{pmatrix} (is/\pi) \right)^4 - \left(\theta \begin{pmatrix} 0 \\ 1 \end{pmatrix} (is/\pi) \right)^4 \right\}, \quad (43)$$

$$Z_\psi^{R-R}(is/\pi) = \frac{1}{(\eta(is/\pi))^4} \left\{ - \left(\theta \begin{pmatrix} 1 \\ 0 \end{pmatrix} (is/\pi) \right)^4 - \left(\theta \begin{pmatrix} 1 \\ 1 \end{pmatrix} (is/\pi) \right)^4 \right\} \quad (44)$$

represent the exchanges of the closed string in NS-NS and R-R modes, respectively. The actual value of this amplitude is zero, which is required by the supersymmetry. The contributions of the NS-NS and R-R exchanges are cancelled out.

The low-energy limit is the $s \rightarrow \infty$ limit in the integrant of the amplitude. In this limit only the exchange of the closed string in lowest-lying states (graviton and dilaton in NS-NS sector) is extracted. It is easy to show that

$$\lim_{s \rightarrow \infty} (\eta(is/\pi))^{-8} Z_\psi^{NS-NS}(is/\pi) = 16 + \mathcal{O}(e^{-2s}), \quad (45)$$

$$\lim_{s \rightarrow \infty} (\eta(is/\pi))^{-8} Z_\psi^{R-R}(is/\pi) = -16 + \mathcal{O}(e^{-2s}), \quad (46)$$

where $\mathcal{O}(e^{-2s})$ is the contribution of massive states. The low-energy limit of the NS-NS contribution of the amplitude is

$$\begin{aligned} A_{NS-NS}^{M^{10}}(y) &= \frac{iV_4}{2\pi (8\pi^2 \alpha')^5} \int_0^\infty ds \left(8\pi^2 \alpha' \frac{\pi}{s}\right)^3 e^{-\frac{y^2}{2\pi\alpha'} \frac{\pi}{s}} \cdot 16 \\ &= iV_4 \cdot 2\pi G_6(y), \end{aligned} \quad (47)$$

where

$$G_d(y) = \frac{1}{4} \pi^{-\frac{d}{2}} \Gamma\left(\frac{d-2}{2}\right) y^{2-d} \quad (48)$$

is the massless scalar Green's function in d dimensions. This amplitude corresponds to the diagram of the tree-level one graviton and one dilaton exchanges between D3-branes in

the supergravity theory with the D-brane effective action of Eq.(1). The amplitude in the supergravity theory can be obtained as

$$A_{NS-NS}^{M^{10}}(y) = iV_4 \cdot 2\kappa^2 \tau_3^2 \cdot G_6(y), \quad (49)$$

where κ is the gravitational constant in ten dimensions. Therefore, we have an expression for D3-brane tension

$$\tau_3^2 = \frac{\pi}{\kappa^2}. \quad (50)$$

The same procedure is possible for general Dp-branes and we have

$$\tau_p^2 = \frac{\pi}{\kappa^2} (4\pi^2 \alpha')^{3-p} \quad (51)$$

which satisfies the T-duality relation

$$\frac{\tau_p}{\tau_{p-1}} = \frac{1}{2\pi\sqrt{\alpha'}}. \quad (52)$$

B. The amplitude in fivebrane backgrounds

The amplitude of Fig.1 in the fivebrane background can be calculated in the same way in the previous subsection.

The transverse dimensions of two parallel D3-branes are $\mu = 1, 2, 3, 4, 5, 6$. The non-trivial gravitational and B-field configurations are realized in the space of $\mu = 7, 8, 9$, namely, the space in D3-brane world-volume. The dilaton background has linear $X^{\mu=6}$ dependence, and the dilaton has a definite (but different) vacuum expectation value in each D3-brane world-volume. The original supersymmetry of $\epsilon_L Q_L + \epsilon_R Q_R$, where Q_L and Q_R are left and right supercharges, is partially broken by this configuration. This fivebrane background in type IIB theory preserves the supersymmetry of

$$\epsilon_L = \Gamma^0 \Gamma^1 \Gamma^2 \Gamma^3 \Gamma^4 \Gamma^5 \epsilon_L, \quad \epsilon_R = -\Gamma^0 \Gamma^1 \Gamma^2 \Gamma^3 \Gamma^4 \Gamma^5 \epsilon_R, \quad (53)$$

and D3-branes preserve the supersymmetry of

$$\epsilon_L = \Gamma^0 \Gamma^7 \Gamma^8 \Gamma^9 \epsilon_R, \quad (54)$$

where Γ^μ is the gamma matrix in ten dimensions (see Ref. [16] for review). Therefore, only eight (1/4) supersymmetries are preserved in this configuration.

Since the open string has no space-time momentum in the direction of $\mu = 1, 2, 3, 4, 5, 6$ and it is stretched between two D3-branes, the terms $\alpha' p^\mu p_\mu$ with $\mu = 1, 2, 3, 4, 5$ and the term $\alpha'(p^{\mu=6} - iQ/\alpha')^2$ for $\mu = 6$ are replaced by a tension term $y^2/4\pi^2\alpha'$ in the world-sheet Hamiltonian, where y is the distance between two parallel D3-branes. Namely, the amplitude is obtained by the replacement

$$iV_7 (8\pi^2 \alpha' t)^{-7/2} \longrightarrow iV_1 (8\pi^2 \alpha' t)^{-1/2} q^{\frac{y^2}{4\pi^2 \alpha'}} \quad (55)$$

in Eq.(35) with $n_{CP}^2 = 2$ and the integration over t :

$$\begin{aligned}
A^{M^6 \otimes W_k^{(4)}} &= \int_0^\infty \frac{dt}{2t} 2 \cdot iV_1 \left(8\pi^2 \alpha' t\right)^{-1/2} e^{-\frac{y^2}{2\pi\alpha'} t} (\eta(it))^{-5} \\
&\times \frac{1}{2} \sum_\delta \sum_{n=0}^k (n+1)(-1)^{\delta n} \chi_k^n(it) \\
&\times \frac{1}{2} \sum_{\alpha,\beta} (-1)^{\alpha+\beta-\alpha\beta} \left(\frac{\theta\left(\frac{\alpha}{\beta}\right)(it)}{\eta(it)} \right)^2 (-1)^{\delta\alpha} \left(\frac{\theta\left(\frac{\alpha}{\beta+\delta}\right)(it)}{\eta(it)} \right)^2.
\end{aligned} \tag{56}$$

To understand this amplitude as the amplitude of a closed string exchange, the valuable t should be changed to $s = \pi/t$. By using Eqs. (40), (41) and (20) we obtain

$$\begin{aligned}
A^{M^6 \otimes W_k^{(4)}} &= \frac{iV_4}{2\pi (8\pi^2 \alpha')^5} \int_0^\infty ds \left(8\pi^2 \alpha' \frac{\pi}{s}\right)^3 e^{-\frac{y^2}{2\pi\alpha'} \frac{\pi}{s}} \\
&\times \frac{(8\pi^2 \alpha')^{3/2} (\eta(is/\pi))^3}{V_3} \cdot \frac{1}{2} \sum_\delta \sum_{n,n'} (n+1)(-1)^{\delta n} S_{nn'} \chi_k^{n'}(is/\pi) \\
&\times (\eta(is/\pi))^{-8} \sum_{\alpha,\beta} (-1)^{\alpha+\beta-\alpha\beta} \left(\frac{\theta\left(\frac{\beta}{\alpha}\right)(is/\pi)}{\eta(is/\pi)} \right)^2 (-1)^{\delta\alpha} \left(\frac{\theta\left(\frac{\beta+\delta}{\alpha}\right)(is/\pi)}{\eta(is/\pi)} \right)^2.
\end{aligned} \tag{57}$$

This amplitude should be compared with the amplitude of Eq.(42). The exchange of the twisted sector ($\delta = 1$) is included in addition to the normal (untwisted) sector ($\delta = 0$).

Now we take the flat space-time limit $k \rightarrow \infty$. First, consider the limit of

$$Z_k^\delta(s) = \sum_{n,n'} (n+1)(-1)^{\delta n} S_{nn'} \chi_k^{n'}(is/\pi). \tag{58}$$

By using Eq.(27) with a condition of $s > s_0 \neq 0$

$$\begin{aligned}
\lim_{k \rightarrow \infty} Z_k^\delta(s) &= \lim_{k \rightarrow \infty} \sum_{n,n'} (n+1)(-1)^{\delta n} \cdot \sqrt{\frac{2}{k+2}} \sin \left(\pi \frac{(n+1)(n'+1)}{k+2} \right) \\
&\times \frac{1}{(\eta(is/\pi))^3} (n'+1) e^{-2(k+2)s \left(\frac{n'+1}{2(k+2)} \right)^2}.
\end{aligned} \tag{59}$$

The summation over n and n' can be carried out in the same way in the previous section, and we obtain

$$\lim_{k \rightarrow \infty} Z_k^\delta(s) = \delta_{\delta 0} \frac{1}{(8\pi^2 \alpha')^{3/2} (\eta(is/\pi))^3} \lim_{k \rightarrow \infty} 2\pi^2 \left(\frac{2}{Q} \right)^3, \tag{60}$$

where $Q = \sqrt{2/(k+2)}$ and we use $\alpha' = 1/2$ in the open string Virasoro character due to the relation between a mode operator and the space-time momentum operator: $\alpha_0^\mu = \sqrt{2\alpha'} p^\mu$. Therefore, the flat space-time limit of the amplitude is

$$\begin{aligned}
\lim_{k \rightarrow \infty} A^{M^6 \otimes W_k^{(4)}} &= \frac{1}{2} \cdot \frac{1}{V_3} \lim_{k \rightarrow \infty} 2\pi^2 \left(\frac{2}{Q} \right)^3 \\
&\times \frac{iV_4}{2\pi (8\pi^2 \alpha')^5} \int_{s_0}^{\infty} ds \left(8\pi^2 \alpha' \frac{\pi}{s} \right)^3 e^{-\frac{y^2}{2\pi \alpha'} \frac{\pi}{s}} \\
&\times (\eta(is/\pi))^{-8} \left\{ Z_{\psi}^{NS-NS}(is/\pi) + Z_{\psi}^{R-R}(is/\pi) \right\}, \tag{61}
\end{aligned}$$

where s_0 is a cut-off by which the contribution from the higher modes of the closed string are eliminated. The first factor $1/2$ is the remnant of the additional GSO projection which realizes the supersymmetry breaking by the fivebrane background. The second factor simply reduces to unity, since $2\pi^2(2/Q)^3$ is the volume of the $SU(2)_k$ manifold of radius $2/Q \rightarrow \infty$ in the $k \rightarrow \infty$ limit. Therefore, the D3-brane tension which is obtained by taking the low-energy limit $s \rightarrow \infty$ in the integrant of Eq.(61) deviates from the one in the flat space-time (Eq.(50)) by a factor of $1/2$:

$$\tau_3^2|_{\text{flat-limit}} = \frac{1}{2} \cdot \frac{\pi}{\kappa^2}. \tag{62}$$

This deviation can be understood as a topological effect which may not disappear in the simple flat space-time limit. Following the discussion in the first section, it is natural to consider that this deviation of the tension suggests the gravitino and/or dilatino condensation by the topological effect.

Here, we give a comment. If s is finite, we can take the flat space-time limit without using the formula of Eq.(20). Namely, we can take the limit in the form of

$$Z_k^{\delta}(\pi/s) = \sum_{n=0}^k (n+1)(-1)^{\delta n} \chi_k^n(i\pi/s). \tag{63}$$

In this case we find that the amplitude, which includes a cut-off $s < s_{\infty}$, also reduces to the one in the flat space-time with an additional factor $1/2$. Therefore, nothing special happens in the region of small s . But, in order to take the flat space-time limit in the way which is consistent with the low-energy limit $s \rightarrow \infty$, we have to use the formula of Eq.(20).

IV. CONCLUSION

The amplitude of the one closed string exchange between two parallel D3-branes in the fivebrane background is calculated. The low-energy limit of the amplitude can be used to obtain the effective D3-brane tension in the low energy supergravity theory. It has been shown that the tension from the flat space-time limit of the amplitude in the fivebrane background does not coincide with the one in the flat space-time. Since the finite curvature effect should disappear in the simple flat space-time limit, the deviation suggests some topological effects.

The effective D3-brane tension in the flat space-time limit is the $1/2$ of the one in the flat space-time. The origin of this factor $1/2$ is the additional GSO projection in the string theory in fivebrane backgrounds, as explained in section II. This additional GSO projection results the spontaneous supersymmetry breaking in fivebrane backgrounds. Therefore, it is

natural to consider that the observed deviation of the effective D-brane tension is related with the supersymmetry breaking.

In the low energy supergravity theory the effective D-brane tension can be changed by the vacuum expectation value of some fields. In case of the curved space-time background the finiteness of the curvature modifies the tension. If the gravitino and/or dilatino condensation by the topological effect occurs as it is expected in Euclidean quantum gravity theory, it also modifies the tension. Since the later modification due to the topological effect should not disappear in the simple flat space-time limit, we can understand that the observed deviation of the effective D-brane tension suggests the gravitino and/or dilatino condensation. This conclusion is supported by the facts that the deviation is related with the supersymmetry breaking and the gravitino condensation triggers supersymmetry breaking.

In this analysis based on the intuitive picture of D-branes it is not clear whether the gravitino condensation really occurs and breaks supersymmetry. The analysis in the language of the boundary states in the world-sheet conformal field theory [17–19] must be important for further investigations. The explicit form of the higher-order terms in the supersymmetric D-brane effective action must be also important to extract the actual values of condensates.

ACKNOWLEDGMENTS

The author would like to thank S. Saito and N. Okada for fruitful discussions and comments.

REFERENCES

- [1] E. Witten, “Dynamical breaking of supersymmetry,” Nucl. Phys. B188 (1981) 513.
- [2] K. Konishi, N. Magnoli and H. Panagopoulos, “Spontaneous breaking of local supersymmetry by gravitational instantons,” Nucl. Phys. B309 (1988) 201; B323 (1989) 441.
- [3] S.-J. Rey, “Confining phase of superstrings and axionic strings,” Phys. Rev. D43 (1991) 526.
- [4] C.G. Callan, J.A. Harvey and A. Strominger, “Worldsheet approach to heterotic instantons and solitons,” Nucl. Phys. B359 (1991) 611.
- [5] I. Antoniadis, S. Ferrara and C. Kounnas, “Exact supersymmetric string solutions in curved gravitational backgrounds,” Nucl. Phys. B421 (1994) 343.
- [6] C. Kounnas, “Type II NS five-branes: non critical strings and their topological sectors,” Fortsch. Phys. 49 (2001) 599.
- [7] J.L. Cardy, “Boundary conditions, fusion rules and the Verlinde formula,” Nucl. Phys. B324 (1989) 581.
- [8] C.P. Bachas, P. Bain and M.B. Green, “Curvature terms in D-brane actions and their M-theory origin,” JHEP 0005 (1999) 011.
- [9] E. Bergshoeff and P.K. Townsend, “Super D-branes,” Nucl. Phys. B490 (1997) 145.
- [10] M. Cederwall, A. von Gussich, B.E.W. Nilsson and A. Westerberg, “The Dirichlet super-three-brane in ten-dimensional type IIB supergravity,” Nucl. Phys. B490 (1997) 163.
- [11] M. Aganagic, C. Popescu and J.H. Schwarz, “Gauge-invariant and gauge-fixed D-brane actions,” Nucl. Phys. B495 (1997) 99.
- [12] P.S. Howe and P.C. West, “The complete $N = 2$, $d = 10$ supergravity,” Nucl. Phys. B238 (1984) 181.
- [13] V.G. Kač and D.H. Peterson, “Infinite-dimensional Lie algebras, theta functions and modular forms,” Adv. Math. 53 (1984) 125.
- [14] D. Gepner and E. Witten, “String theory on group manifold,” Nucl. Phys. B278 (1986) 493.
- [15] J. Polchinski, “String theory,” Vol. I and II, Cambridge Univ. Press (1998).
- [16] A. Giveon and D. Kutasov, “Brane dynamics and gauge theory,” Rev. Mod. Phys. 71 (1999) 983.
- [17] A. Recknagel and V. Schomerus, “D-branes in Gepner models,” Nucl. Phys. B531 (1998) 185.
- [18] J. Maldacena, G. Moore and N. Seiberg, “Geometrical interpretation of D-branes in gauged WZW models,” JHEP 0107 (2001) 046.
- [19] J. Maldacena, G. Moore and N. Seiberg, “D-brane Charges in Five-brane backgrounds,” hep-th/0108152.

FIGURES




FIG. 1. The diagram of one closed string exchange between two parallel D3-branes